

1982


RADIOCARBON DATING IN A KARSTIC COASTAL AQUIFER IN TROPICAL CLIMATIC CONDITIONS.

D.M. ARELLANO Ph.D., J.R. FAGUNDO Ph.D., J. ŠILAR Ph.D., P. JILEK (CUBA)

The results of the determination of both Calcite Saturation Ratio and PCO_2 , they measured in seven wells distributed in an area of karstic coastal aquifer, are presented. The ^{14}C activity and the groundwater age were calculated by using of four models reported in consulted bibliography, [1,2].

The intrusion of water from the sea, into the coastal aquifer, was examined by means of the Specific Conductivity, Total Dissolved Solids, Ionic Ratios (Mg^{2+}/Ca^{2+} , Cl^-/HCO_3^- , Na^+/Ca^{2+} , among others). This intrusion determines the kinetic factors of the water-geologic medium system.

The intrusion effect on the limestones produces the oversaturation of the dissolution; it contributes in the obtention of low activity of ^{14}C , and, consequently, "ages" which range from 1000 up to 16 000 years are obtained. These "ages" are not in agreement with the available preliminary information related to the isotopic composition ($\delta^{18}O$, δ^2H) measured in rainfalls, springs and wells and the 3H measurements [3] which indicates the occurrence of recent waters.



The effect of the subsurface karstic cavities and fissures influences the low ^{14}C activity calculated. In this case, the diffusivity of the rocks affect the activity measured, according to the teorethical formulations and investigations made in the fissured medium with porous matrix [4,5].

The conditions of the humid tropical climate are shown. In these conditions, the biological activity is intensive and the PCO_2 values calculated are in an order from 10^{-2} - 10^{-3} atm.

In this karstic confined aquifer, the transmissivity values range from 5 000 to 20 000 $\text{m}^2 \cdot \text{day}^{-1}$ and the discharge at the wells, during the pumping test, can reach up to 390 l.s^{-1}

REFERENCES

- 1 GEYH, M.A., Basic studies in hydrology and ^{14}C and ^3H measurements.
International Geolog. Congress, 24th Sess.
Sect. 11 Hydrogeology, Canada (1972) 227-234.
- 2 SALEM, O. et al, Groundwater flow patterns in the western Lybian Arab. Jamahiriya evaluated from isotopic data, Arid Zone Hydrology Investigations with isotope techniques, Proceed. Panel IAEA, Vienna (1980) 165-179.
- 3 ARELLANO, D.M., SEILER, K.P, RAUERT, W., Estudio Isotópico de la llanura costera sur de Pinar del Río, Programa OIEA/GSF (in preparation).
- 4 NERETNICKS, I., Diffusion in the rock matrix, an important factor in radionuclide retardation?, J. Geophys. Research 85 B8, (1980) 4379-81.
- 5 MALOSZEWSKI, P., ZUBER, A. Interpretation of artificial and environmental tracers in fissured rocks with a porous matrix. Report No. 1221/AP. Institute of Nuclear Physics, Krakow, Poland (1983) 1-17.