

Cretaceous-Paleogene boundary deposits at Loma Capiro, central Cuba: Evidence for the Chicxulub impact

Laia Alegret* Department of Earth Sciences, University College London, WC1E 6BT London, UK, and Departamento de Ciencias de la Tierra, Universidad de Zaragoza, 50009 Zaragoza, Spain

Ignacio Arenillas }
José A. Arz } Departamento de Ciencias de la Tierra, Universidad de Zaragoza, 50009 Zaragoza, Spain

Consuelo Díaz Instituto de Geología y Paleontología, La Habana, Cuba

José M. Grajales-Nishimura Instituto Mexicano del Petróleo, 07730 Mexico D.F., Mexico

Alfonso Meléndez }
Eustoquio Molina } Departamento de Ciencias de la Tierra, Universidad de Zaragoza, 50009 Zaragoza, Spain

Reinaldo Rojas Museo Nacional de Historia Natural, 10100 La Habana, Cuba

Ana R. Soria Departamento de Ciencias de la Tierra, Universidad de Zaragoza, 50009 Zaragoza, Spain

ABSTRACT

A newly discovered Upper Cretaceous to lower Paleogene section at Loma Capiro (central Cuba) has provided new evidence for a Cretaceous-Paleogene boundary age for the Chicxulub impact. The studied sediments at Loma Capiro consist of a foraminifera-rich marl and sandstone hemipelagic sequence, and a 9.6-m-thick intercalated clastic complex. Planktic foraminifera indicate an upper Maastrichtian age for the sediments below the clastic complex and a lowermost Danian age for those just above this complex. Small benthic foraminifera from below and above the clastic complex indicate deposition at middle to lower bathyal depths. The fining-upward clastic complex consists of a basal breccia that is overlain by microconglomerates and coarse- to fine-grained sandstones. The clastic complex contains reworked foraminifera from different ages and different paleoenvironments and, toward the top, impact material such as altered microtektites, shocked quartz, terrestrial chondrules, and accretionary lapilli. These microfacies suggest deposition from gravity flows that eroded sediments from upper-slope and shelf settings and redeposited them in deeper bathyal environments. We suggest that the origin of the clastic complex may be linked to the collapse of the Cuban platform, triggered by the Cretaceous-Paleogene impact at Chicxulub.

Keywords: Chicxulub impact, Cretaceous-Paleogene event, gravity flows, foraminifera, Cuba.

INTRODUCTION

The Cretaceous-Paleogene boundary event and the related catastrophic mass extinction are usually explained as the result of a large asteroid impact that occurred at Chicxulub in the Yucatan Peninsula, Mexico. The impact scenario is broadly accepted to account for the origin of the Cretaceous-Paleogene clastic deposits from the Gulf of Mexico, Caribbean, and North Atlantic (Smit et al., 1996; Grajales-Nishimura et al., 2000; Klaus et al., 2000; Soria et al., 2001; Tada et al., 2003). The Chicxulub impact caused the destabilization of the central and North American continental margins and megatsunamis, triggering the deposition of the so-called “Cretaceous-Paleogene boundary cocktail” unit in these areas (Bralower et al., 1998). This unit includes a mixture of reworked microfossils from different ages, impact-derived materials, and heterogeneous lithic fragments.

This single catastrophic scenario is not supported by a small group of scientists who argue that the Chicxulub crater was formed 300

k.y. earlier. This hypothesis is based on two main arguments. The first is the presence of several horizons of spherules found in the La Sierrita area (northeastern Mexico; Stinnesbeck et al., 2001), although the apparent multiple spherule horizons may be explained by slumping (Soria et al., 2001). The second is the presence of a 50-cm-thick layer of micritic limestone containing uppermost Maastrichtian planktic foraminiferal assemblages overlying the impact breccia in the Yaxcopoil-1 drill hole (Chicxulub crater; Keller et al., 2004). However, such a critical interval is not a micritic limestone, but it consists of a dolomitic calcareous sandstone with climbing ripples and scarce, reworked Albian to Maastrichtian

planktic foraminifera, and it probably represents the infill of the crater due to the marine invasion into the cavity that occurred just after the Cretaceous-Paleogene boundary impact (Arz et al., 2004; Smit et al., 2004).

The origin and model of deposition of the Cretaceous-Paleogene boundary deposits from the Gulf Coast and Caribbean are controversial issues that need more research. Studies performed in western Cuba have documented the presence of a clastic megabed, the origin of which has been related to the Chicxulub impact (e.g., Tada et al., 2003). Studies of Cretaceous-Paleogene boundary deposits from central Cuba are scarce, and the paleobathymetry and dating of these sediments, as well as the environmental consequences of such an impact, are not yet clear. In order to clarify these issues, we performed a multidisciplinary study of the new Loma Capiro section (central Cuba), including detailed stratigraphic, petrographic, and paleobathymetric analyses, and high-resolution foraminiferal biostratigraphy. In contrast to other Cretaceous-Paleogene boundary clastic deposits in sections and wells from Mexico, Guatemala, Belize, or from the crater itself, the newly discovered Loma Capiro section contains abundant microfossils, which provide some additional information on the origin and deposition of this unit, and on its relationship with the Chicxulub impact and the Cretaceous-Paleogene boundary event.

MATERIAL AND METHODS

The Loma Capiro section is located in central Cuba (Villa Clara Province; Fig. 1), on the




Figure 1. Location of Loma Capiro section.

*Corresponding author e-mail: laia@unizar.es.




Figure 3. A: Slumps affecting Cretaceous sediments at Loma Capiro. B: Calcified broken vesiculated spherule. C: Shocked quartz grain with one set of planar deformation features. D: Accretionary lapilli, showing concentric structures and core lined by dark rim, and outer dark gray band.

spherules have been altered to calcite; they are spherical and normally vesicular.

The clastic complex is overlain by a Danian sequence of tabular bodies, as much as 2 m thick, of ochre and gray massive to occasionally laminated silts, intercalated with 0.25–0.8-m-thick tabular sandstone levels. The sandstones are fine grained and display parallel lamination and low-angle cross-lamination.

BIOSTRATIGRAPHY

For biostratigraphic determinations (Fig. 2), we used the planktic foraminiferal zonations by Arz and Molina (2002) and Arenillas et al. (2004). We recognized 67 planktic foraminiferal species in the lowermost 6 m of the section (Cretaceous autochthonous sediments not affected by slumping), which belong to the *Pseudoguembelina hariaensis* subzone (*Abathomphalus mayaroensis* zone). The uppermost Maastrichtian *Plummerita hantkeninoides* subzone, and probably most of the *Pseudoguembelina hariaensis* subzone, are absent. The coexistence of *P. hariaensis*, *Globotruncana bulloides*, and *Contusotruncana formicata* in this interval indicates a stratigraphic gap that affects the last ~1500 k.y. of the upper Maastrichtian (Arz and Molina, 2002).

Above the clastic complex, the *Guembelitra cretacea*, *Parvularugoglobigerina* (*Pv.*) *eugubina*, *Parasubbotina pseudobulloides*, and *Globanomalina compressa* zones were

identified. Since the clastic complex only contains reworked microfossils, we have included it in a barren interzone between the *Abathomphalus mayaroensis* zone and the *Guembelitra cretacea* zone. A short hiatus comprising the *Hedbergella holmdelensis* subzone (= P0 by Beggren et al., 1995) and most of the *Pv. longiapertura* subzone affects the lowermost part of the Paleocene (i.e., the first 15–20 k.y. of the Danian; Arenillas et al., 2004). Therefore, the *G. cretacea* zone is practically absent at Loma Capiro, and it is only represented by 4 cm identified between the clastic complex (barren interzone) and the *Pv. eugubina* zone.

PALEOBATHYMETRY

Benthic foraminiferal assemblages at Loma Capiro contain abundant representatives of the bathyal to abyssal Velasco-type fauna, such as *Aragonia velascoensis*, *Gaudryina pyramidata*, *Gyroidinoides globosus*, *Nuttallides truempyi*, *Nuttallinella florealis*, or *Stensioeina beccariiiformis*, as well as other taxa typical of middle bathyal to abyssal depths, namely *A. ouezzanensis*, *Gyroidinoides* spp., *Marssonella oxycona*, *Nt. coronula*, *Oridorsalis umbonatus*, *Paralabamina hillebrandti*, and *Spiroplectammina spectabilis* (e.g., Alegret et al., 2001, 2003). Species with an upper depth limit of ~500–700 m (e.g., *Bulimina trinitatensis*, *N. truempyi*, *Nt. florealis*, *S. beccariiiformis*, *Sp. spectabilis*; Alegret et al., 2003) are most abundant at Loma Capiro. These data indicate that sediments below (Maastrichtian)

and above (Danian) the clastic complex were deposited at middle to lower bathyal depths.

AGE AND EMPLACEMENT OF THE CLASTIC COMPLEX

Abundant reworked foraminifera, including nearly all planktic foraminiferal species identified in the Maastrichtian marls, were found in the clastic complex. Specimens of *Globotruncana linneiana* and *Contusotruncana plummerae* (upper Santonian–lower Maastrichtian), as well as *P. hantkeninoides* and *Ps. hariaensis* (uppermost Maastrichtian), were also found. These species do not coincide in age, suggesting the presence of reworked mixed Upper Cretaceous assemblages in this unit. The presence of reworked specimens of *P. hantkeninoides* in the clastic complex at Loma Capiro indicates that the stratigraphic interval that belonged to the *P. hantkeninoides* subzone (i.e., the uppermost 300 k.y. of the Maastrichtian; Keller et al., 2004) was present in this area, and that it was probably eroded during the Cretaceous–Paleogene event. According to these data, the Chicxulub impact event cannot have taken place 300 k.y. before the Cretaceous–Paleogene boundary, as suggested by Stinnesbeck et al. (2001) and Keller et al. (2004). On the contrary, these data strongly support the Cretaceous–Paleogene boundary age of the Chicxulub impact.

Accordingly, benthic macroforaminiferal assemblages from the clastic complex consist of a mixture of species from different ages, such as *Asterorbis aguayoi* (Campanian–lower Maastrichtian), *A. macei*, *A. rooki*, *Pseudorbitoides trechmani*, *Sulcoperculina globosa*, *Torreina torrei* (Maastrichtian), and *A. havanensis*, *A. cubensis*, and *Orbitoides apiculata apiculata* (upper Maastrichtian).

The clastic complex contains a mixture of deep-water benthic foraminiferal species (e.g., *A. velascoensis*, *B. trinitatensis*, *G. pyramidata*, *Gy. globosus*, *N. truempyi*, *Nt. coronula*, *Nt. florealis*, *P. hillebrandti*, *Osangularia velascoensis*, *S. beccariiiformis*) and shallow-water species (e.g., *Anomalinoidea praeacutus*, *Coryphostoma incrassata* forma *gigantea*, *Oridorsalis plummerae*) that are common in sublittoral to upper bathyal environments (e.g., Alegret et al., 2003). Moreover, assemblages from the second and third subunits contain bryozoans and abundant macroforaminifera from sublittoral environments. These data suggest that allochthonous Upper Cretaceous material coming from the shelf and the upper part of the slope was transported and redeposited at middle-lower bathyal depths during the deposition of the clastic complex.

Diverse impact material was found toward the top of the clastic complex. The chondrule-like particles and the accretionary lapilli are similar to those reported from the Miocene Ries Crater in Germany and from the Cretaceous–

Paleogene boundary El Guayal section in Mexico, which were interpreted as having been generated by impact mechanisms (Graup, 1981; Grajales-Nishimura et al., 2003). The calcified glass spherules have been interpreted as altered tektites and have been reported from many Cretaceous-Paleogene boundary localities in Mexico (e.g., Smit et al., 1996). Altered microtektites have also been documented from the Cretaceous-Paleogene fireball layer in North America, Canada, the Atlantic Ocean, Southern Ocean, and Mediterranean Domain, as well as in Boreal Paratethys (Ortega-Huertas et al., 2002).

The aforementioned characteristics of the clastic complex indicate that it is equivalent to the Cretaceous-Paleogene boundary cocktail unit defined by Bralower et al. (1998), which has been identified in sections and wells from the Chicxulub crater and the Gulf Coast margins, and is commonly related to the Chicxulub impact event (Smit et al., 1996; Grajales-Nishimura et al., 2000; Arz et al., 2004).

The sedimentologic features and the micro-paleontologic and mineralogic content of the Cretaceous-Paleogene clastic complex allowed us to infer that it corresponds to a mass flow, which includes (1) coarse-grained debris flows that led to the deposition of the breccia, which also contains clasts from multiple sources; and (2) slumping processes that locally affected both the breccia and the underlying Cretaceous marly sediments. Similar coarse breccia units have been reported from around the Chicxulub impact crater (Grajales-Nishimura et al., 2003; Tada et al., 2003), and have been related to the collapse of the Mexican and Cuban platforms, and to the large-scale gravity flows triggered by the seismic waves originated by the Cretaceous-Paleogene impact at Chicxulub.

We interpret the upper subunit as due to the settling of a resuspended sediment cloud produced by the entrainment into the basin of fine material from the mass flow. This model also accounts for the concentration of impact evidence toward the top of the clastic complex. Alternatively, the upper subunit might correspond to the homogenite unit documented by Tada et al. (2003) from western Cuba. According to their model, the homogenite unit was deposited in a few weeks after the Chicxulub impact, from a resuspended sediment cloud that was generated in shallower areas by large tsunamis.

CONCLUSIONS

Sedimentological features and paleobathymetrical data inferred from benthic foraminifera from below and above the clastic complex indicate hemipelagic sedimentation at middle to lower bathyal depths. Biostratigraphic in-

ferences based on planktic foraminifera indicate a late Maastrichtian age (*Pseudoguembelina hariaensis* subzone) for the autochthonous sediments below the clastic complex, and an earliest Paleogene age (*Guembelitra cretacea* to *Globanomalina compressa* zones) for those just above this complex.

The presence of impact material and reworked uppermost Maastrichtian foraminifera in the clastic complex supports the relationship between the Chicxulub impact and the Cretaceous-Paleogene boundary event. We suggest that the uppermost Maastrichtian gap was caused by erosion associated with the gravity flows and slumping processes that originated during the Cretaceous-Paleogene impact event. We conclude that the deposition of the clastic complex was geologically instantaneous and linked to a rapid and catastrophic event (Chicxulub impact) at the Cretaceous-Paleogene boundary.

ACKNOWLEDGMENTS

We thank Chris Kilburn (University College London) and reviewers T.J. Bralower and B.T. Huber for valuable suggestions that significantly improved our paper. Research was funded by Projects CGL2004-00738/BTE (Spain), IMP-D 01003 (Mexico), and PNAP 035 (Cuba). L. Alegret holds a postdoctoral grant (EX2003-0007).

REFERENCES CITED

- Alegret, L., Molina, E., and Thomas, E., 2001, Benthic foraminifera at the Cretaceous-Tertiary boundary around the Gulf of Mexico: *Geology*, v. 29, p. 891–894, doi: 10.1130/0091-7613(2001)029<0891:BFATCT>2.0.CO;2.
- Alegret, L., Molina, E., and Thomas, E., 2003, Benthic foraminiferal turnover across the Cretaceous/Tertiary boundary at Agost (southeastern Spain): Paleoenvironmental inferences: *Marine Micropaleontology*, v. 48, p. 251–279.
- Arenillas, I., Arz, J.A., and Molina, E., 2004, A new high-resolution planktic foraminiferal zonation and subzonation for the lower Danian: *Lethaia*, v. 37, p. 79–95, doi: 10.1080/00241160310005097.
- Arz, J.A., and Molina, E., 2002, Biostratigrafía y cronostratigrafía con foraminíferos planctónicos del Campaniense superior y Maastrichtiense de latitudes subtropicales y templadas (España, Francia y Tunicia): *Neues Jahrbuch für Geologie und Paläontologie Abhandlungen*, v. 224, p. 161–195.
- Arz, J.A., Alegret, L., and Arenillas, I., 2004, Foraminiferal biostratigraphy and paleoenvironmental reconstruction at Yaxcopoil-1 drill hole (Chicxulub crater, Yucatan Peninsula): *Meteoritics & Planetary Science*, v. 39, p. 1099–1111.
- Berggren, W.A., Kent, D.V., Swisher, C.C., III, and Aubry, M.P., 1995, A revised Paleogene geochronology and chronostratigraphy, in Berggren, W.A., et al., eds., *Geochronology, time scales and global stratigraphic correlation: Society for Sedimentary Geology (SEPM) Special Publication 54*, p. 129–213.
- Bralower, T.J., Paull, C.K., and Leckie, R.M., 1998, The Cretaceous-Tertiary boundary cocktail: Chicxulub impact triggers margin collapse and extensive sediment gravity flows: *Geology*, v. 26, p. 331–334, doi: 10.1130/0091-7613(1998)026<0331:TCTBCC>2.3.CO;2.
- Grajales-Nishimura, J.M., Cedillo, E., Rosales, M.C., Morán, D.J., Alvarez, W., Claes, P., Ruiz, J., García, J., Padilla, P., and Sánchez, A., 2000, Chi-

cxulub impact: The origin of reservoir and seal facies in the southeastern Mexico oil fields: *Geology*, v. 28, p. 307–310, doi: 10.1130/0091-7613(2000)028<0307:CITOOOR>2.3.CO;2.

- Grajales-Nishimura, J.M., Murillo, G., Rosales, C., Cedillo, E., and García, J., 2003, Heterogeneity of lithoclast composition in the deep-water carbonate breccias of the K/T Campeche, in Bartolini, C., et al., eds., *The circum-Gulf of Mexico and the Caribbean: Hydrocarbon habitats, basin formation, and plate tectonics: American Association of Petroleum Geologists Memoir 79*, p. 312–329.
- Graup, G., 1981, Terrestrial chondrules, glass spherules, and accretionary lapilli from the suevite, Ries Crater, Germany: *Earth and Planetary Science Letters*, v. 55, p. 407–418, doi: 10.1016/0012-821X(81)90168-0.
- Keller, G., Adatte, T., Stinnesbeck, W., Stüben, D., Berner, Z., Kramar, U., and Harting, M., 2004, More evidence that the Chicxulub impact predates the K/T mass extinction: *Meteoritics & Planetary Science*, v. 39, p. 1127–1144.
- Klaus, A., Norris, R.D., Kroon, D., and Smit, J., 2000, Impact-induced mass wasting at the KT boundary: Blake Nose, western North Atlantic: *Geology*, v. 28, p. 319–322, doi: 10.1130/0091-7613(2000)028<0319:IIMWAT>2.3.CO;2.
- Ortega-Huertas, M., Martínez-Ruiz, F., Palomo, I., and Chamley, H., 2002, Review of the mineralogy of the Cretaceous-Tertiary boundary clay: Evidence supporting a major extraterrestrial catastrophic event: *Clay Minerals*, v. 37, p. 395–411, doi: 10.1180/0009855023730054.
- Smit, J., Roep, T.B., Alvarez, W., Montanari, A., Claes, P., Grajales-Nishimura, J.M., and Bermudez, J., 1996, Coarse-grained, clastic sandstone complex at the K/T boundary around the Gulf of Mexico: Deposition by tsunami waves induced by the Chicxulub impact?, in Ryder, G., et al., eds., *The Cretaceous-Tertiary event and other catastrophes in Earth history: Geological Society of America Special Paper 307*, p. 151–182.
- Smit, J., van der Gaast, S., and Lustenhouwer, W., 2004, Is the transition impact to post-impact rock complete? Some remarks based on XFR scanning, electron microprobe, and thin section analyses of the Yaxcopoil-1 core in the Chicxulub crater: *Meteoritics & Planetary Science*, v. 39, p. 1113–1126.
- Soria, A.R., Liesa, C., Mata, M.P., Arz, J.A., Alegret, L., Arenillas, I., and Meléndez, A., 2001, Slumping and a sandbar deposit at the K/T boundary in the El Tecolote section (northeastern Mexico): An impact-induced sediment gravity flow: *Geology*, v. 29, p. 231–234, doi: 10.1130/0091-7613(2001)029<0231:SAASDA>2.0.CO;2.
- Stinnesbeck, W., Schulte, P., Lindenmaier, F., Adatte, T., Affolter, M., Schilli, L., Keller, G., Stüben, D., Berner, Z., Kramar, U., Burns, S.J., and López-Oliva, J.G., 2001, Late Maastrichtian age of spherule deposits in northeastern Mexico: Implication for Chicxulub scenario: *Canadian Journal of Earth Sciences*, v. 38, p. 229–238, doi: 10.1139/cjes-38-2-229.
- Tada, R., Iturralde, M.A., Matsui, T., Tajika, E., Oji, T., Goto, K., Nakano, Y., Takayama, H., Yamamoto, S., Kiyokawa, S., Toyoda, K., García, D., Díaz, C., and Rojas, R., 2003, K/T boundary deposits in the paleo-western Caribbean basin, in Bartolini, C., et al., eds., *The circum-Gulf of Mexico and the Caribbean: Hydrocarbon habitats, basin formation, and plate tectonics: American Association of Petroleum Geologists Memoir 79*, p. 582–604.

Manuscript received 1 February 2005

Revised manuscript received 13 May 2005

Manuscript accepted 16 May 2005

Printed in USA